

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

## AD-A275 286



tion is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this burden, including this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Avenue, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

2. REPORT DATE  
1993

3. REPORT TYPE AND DATES COVERED  
Interim 92/01/01 - 92/12/31

4. TITLE AND SUBTITLE

Distributional records for Aedes mosquitoes from the U.S. Air Force ovitrapping program - 1992

5. FUNDING NUMBERS

6. AUTHOR(S)

Chad P. McHugh

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Armstrong Laboratory  
Occupational and Environmental Health Directorate  
Brooks Air Force Base, TX 78235-5114

8. PERFORMING ORGANIZATION REPORT NUMBER

AL-JA-1993-0002

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)

Armstrong Laboratory  
Occupational and Environmental Health Directorate  
Brooks Air Force Base, TX 78235-5114

10. SPONSORING / MONITORING AGENCY REPORT NUMBER

11. SUPPLEMENTARY NOTES

To be published in the Journal of the American Mosquito Control Association

12a. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for public release; distribution is unlimited

12b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 words)

During 1992, ovitrapping to sample container-breeding Aedes was conducted at 28 U.S. Air Force installations in North America. Aedes albopictus was collected at 19 installations. The collection of Ae. albopictus at Goodfellow Air Force base (AFB), TX and Moody AFB, GA, represent new records for Tom Green and Lowndes counties, respectively. Six installations were positive for Ae. aegypti; 13 were positive for Ae. triseriatus.

DTIC  
SELECTED  
FEB 01 1994  
S B D

14. SUBJECT TERMS

ovitrapping oviposition mosquito Aedes aegypti  
Aedes albopictus Aedes triseriatus distribution

15. NUMBER OF PAGES

4

16. PRICE CODE

17. SECURITY CLASSIFICATION OF REPORT

UNCLASSIFIED

18. SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED

19. SECURITY CLASSIFICATION OF ABSTRACT

UNCLASSIFIED

20. LIMITATION OF ABSTRACT

UL

## DISTRIBUTIONAL RECORDS FOR *Aedes* MOSQUITOES FROM THE U.S. AIR FORCE OVITRAPPING PROGRAM—1992

CHAD P. McHUGH<sup>1</sup>

Occupational and Environmental Health Directorate  
Armstrong Laboratory, Brooks Air Force Base, TX 78235

**ABSTRACT.** During 1992, ovitrapping to sample populations of container-breeding *Aedes* was conducted at 28 U.S. Air Force installations in North America. *Aedes albopictus* was collected at 19 installations. The collections of *Ae. albopictus* at Goodfellow Air Force base (AFB), TX, and Moody AFB, GA, represent new records for Tom Green and Lowndes counties, respectively. Six installations were positive for *Ae. aegypti*; 13 were positive for *Aedes triseriatus*.

U.S. Air Force (USAF) regulations mandate the use of oviposition traps to monitor container-breeding *Aedes* at installations in the known, or potential, range of the Asian tiger mosquito, *Aedes albopictus* (Skuse). This sampling was dictated by concern about the potential for transmission of dengue or yellow fever viruses should these exotic viruses be introduced into the USA. Additionally, the recent isolation of eastern equine encephalitis virus from *Ae. albopictus* in Florida (Mitchell et al. 1992) emphasizes the possibility that this mosquito may serve as a vector of viruses already enzootic in this country.

During 1992, 28 USAF installations submitted a total of 1,608 ovipaddles to the Occupational Medicine Division, Brooks AFB for processing (Table 1). Eggs on ovipaddles were tentatively identified as *Aedes triseriatus* (Say) or *Aedes (Stegomyia)* sp. based on their color and surface texture (Linley 1989a, 1989b) and held for a few days at high humidity to ensure embryonation. The eggs were then allowed to dry and were subsequently hatched in a 1:1 mixture of tap and distilled water. Larvae were fed liver powder and reared to the 4th instar or adult stage for specific identification. Records of previous years' collections at USAF installations have been reported by McHugh and Vande Berg (1989), McHugh and Hanny (1990), McHugh (1991) and McHugh (1992).

Nineteen installations submitted ovipaddles positive for *Ae. albopictus* (Table 1). At these bases, the prevalence was 30.7% (401/1,308). If the 139 *Aedes (Stegomyia)*-positive ovipaddles from these 19 bases are assumed to be *Ae. albopictus*, the prevalence at positive bases would be 41.3% (540/1,308).

The collection of *Ae. albopictus* at Moody AFB, GA, on August 31, is believed to be a first record

for Lowndes County. Additional *Ae. albopictus*- and (*Stegomyia*)-positive ovipaddles were collected through September 30 when sampling stopped. *Aedes albopictus* is widespread in nearby northern Florida (O'Meara et al. 1992), and was subsequently collected in October in suburban Valdosta, GA, about 7 km south of Moody AFB (M. L. Womack, unpublished data).

The collection of *Ae. albopictus* at Goodfellow AFB, TX, on July 8 is the first record of this species in Tom Green County. An ovipaddle with eggs tentatively identified as *Aedes (Stegomyia)* sp. was collected on June 30, but specimens could not be reared for specific identification. Additional *Ae. albopictus*- and (*Stegomyia*)-positive ovipaddles were collected through August 27. Sampling at Goodfellow AFB ended September 23. Because there has been only limited sampling at Goodfellow in the previous 4 years (5 ovipaddles, all negative, submitted during 1989), it cannot be determined when the species was introduced into this area.

The trend at Tinker AFB, OK, is perhaps typical of installations that *Ae. albopictus* has recently colonized. The Asian tiger mosquito was first collected at that base in 1990 when 4.4% of ovipaddles were positive (McHugh 1991). This prevalence increased to 25.6 and 45.8% in 1991 (McHugh 1992) and 1992, respectively. If *Aedes (Stegomyia)*-positive ovipaddles are included (*Aedes aegypti* (Linn.) has not been collected at Tinker AFB), these figures become 8.8, 32.6 and 54.1%.

*Aedes aegypti* was collected at 6 installations. This species has been present in low numbers (<7%) at Seymour Johnson AFB, NC, for 3 of the previous 4 years in the absence of *Ae. albopictus*. Sampling at Shaw AFB, SC, has detected *Ae. aegypti* sporadically, including 1991 when *Ae. albopictus* was first collected at that base and again in 1992. Three bases in San Antonio, TX—Brooks AFB, Lackland AFB, and Kelly AFB—recorded the presence of *Ae. aegypti*, but

<sup>1</sup> Mail address: AL/OEMB (Attn: C. P. McHugh), 2402 E. Drive, Brooks AFB TX 78235-5114.



Table 1. Summary of ovipaddles processed at the Occupational and Environmental Health Directorate, Armstrong Laboratory, Brooks Air Force Base, TX, during 1992.

| Installation <sup>1</sup> and state | County<br>(Parish in LA) | Total<br>ovipaddles | Positive ovitraps     |      |                    |     |                                    |      |     |      |     |      |
|-------------------------------------|--------------------------|---------------------|-----------------------|------|--------------------|-----|------------------------------------|------|-----|------|-----|------|
|                                     |                          |                     | <i>Ae. albopictus</i> |      | <i>Ae. aegypti</i> |     | <i>Ae. (Stegomyia) triseriatus</i> |      |     |      |     |      |
|                                     |                          |                     | n                     | %    | n                  | %   | n                                  | %    | n   | %    | n   | %    |
| Gunter AFB, AL                      | Montgomery               | 8                   | 5                     | 62.5 | 0                  | 0.0 | 2                                  | 25.0 | 0   | 0.0  | 0   | 0.0  |
| Maxwell AFB, AL                     | Montgomery               | 19                  | 11                    | 57.9 | 0                  | 0.0 | 1                                  | 5.3  | 6   | 31.6 | 6   | 31.6 |
| Little Rock AFB, AR                 | Pulaski                  | 164                 | 47                    | 28.7 | 0                  | 0.0 | 15                                 | 9.1  | 15  | 9.1  | 15  | 9.1  |
| Bolling AFB, DC                     | —                        | 40                  | 0                     | 0.0  | 0                  | 0.0 | 0                                  | 0.0  | 4   | 10.0 | 4   | 10.0 |
| Avon Park AFR, FL                   | Polk                     | 59                  | 5                     | 8.5  | 0                  | 0.0 | 4                                  | 6.8  | 0   | 0.0  | 0   | 0.0  |
| MacDill AFB, FL                     | Hillsborough             | 40                  | 6                     | 15.0 | 0                  | 0.0 | 1                                  | 2.5  | 0   | 0.0  | 0   | 0.0  |
| Tyndall AFB, FL                     | Bay                      | 20                  | 1                     | 5.0  | 0                  | 0.0 | 0                                  | 0.0  | 0   | 0.0  | 0   | 0.0  |
| Moody AFB, GA                       | Lowndes                  | UNK <sup>2</sup>    | P                     | —    | N                  | —   | P                                  | —    | N   | —    | N   | —    |
| Barksdale AFB, LA                   | Bossier                  | 57                  | 24                    | 42.1 | 0                  | 0.0 | 5                                  | 8.8  | 4   | 7.0  | 4   | 7.0  |
| Columbus AFB, MS                    | Lowndes                  | 85                  | 32                    | 37.6 | 0                  | 0.0 | 14                                 | 16.5 | 31  | 36.5 | 31  | 36.5 |
| Keesler AFB, MS                     | Harrison                 | 10                  | 9                     | 90.0 | 0                  | 0.0 | 1                                  | 10.0 | 0   | 0.0  | 0   | 0.0  |
| Pope AFB, NC                        | Cumberland               | 22                  | 0                     | 0.0  | 0                  | 0.0 | 0                                  | 0.0  | 1   | 4.5  | 1   | 4.5  |
| Seymour Johnson AFB, NC             | Wayne                    | 142                 | 0                     | 0.0  | 5                  | 3.5 | 4                                  | 2.8  | 1   | 0.7  | 1   | 0.7  |
| Tinker AFB, OK                      | Oklahoma                 | 24                  | 11                    | 45.8 | 0                  | 0.0 | 2                                  | 8.3  | 0   | 0.0  | 0   | 0.0  |
| Charleston AFB, SC                  | Berkeley                 | 58                  | 3                     | 5.2  | 0                  | 0.0 | 0                                  | 0.0  | 1   | 1.7  | 1   | 1.7  |
| Shaw AFB, SC                        | Sumter                   | UNK <sup>2</sup>    | P                     | —    | P                  | —   | P                                  | —    | P   | —    | P   | —    |
| Arnold AFS, TN                      | Coffee                   | 55                  | 25                    | 45.5 | 0                  | 0.0 | 4                                  | 7.3  | 29  | 52.7 | 29  | 52.7 |
| Brooks AFB, TX                      | Bexar                    | 130                 | 66                    | 50.8 | 1                  | 0.8 | 15                                 | 11.5 | 1   | 0.8  | 1   | 0.8  |
| Dyess AFB, TX                       | Taylor                   | 3                   | 0                     | 0.0  | 0                  | 0.0 | 1                                  | 33.3 | 0   | 0.0  | 0   | 0.0  |
| Goodfellow AFB, TX                  | Tom Green                | 111                 | 4                     | 3.6  | 1                  | 0.9 | 6                                  | 5.4  | 0   | 0.0  | 0   | 0.0  |
| Kelly AFB, TX                       | Bexar                    | 79                  | 26                    | 32.9 | 1                  | 1.3 | 4                                  | 5.1  | 1   | 1.3  | 1   | 1.3  |
| Lackland AFB, TX                    | Bexar                    | 225                 | 76                    | 33.8 | 2                  | 0.9 | 51                                 | 22.7 | 8   | 3.6  | 8   | 3.6  |
| Randolph AFB, TX                    | Bexar                    | 164                 | 50                    | 30.5 | 0                  | 0.0 | 14                                 | 8.5  | 0   | 0.0  | 0   | 0.0  |
| Total                               |                          | 1,608 <sup>3</sup>  | 401                   | 24.9 | 10                 | 0.6 | 144                                | 8.9  | 102 | 6.3  | 102 | 6.3  |

<sup>1</sup> Installation abbreviations: Air Force Base, AFB; Air Force Station, AFS; Space Center, SC.

<sup>2</sup> The total number of ovitraps used was unknown (UNK). The installation was positive (P) or negative (N) for the species indicated.

<sup>3</sup> Includes ovipaddles submitted by the following installations, all of which were negative: Homestead AFB, FL—31; Kennedy SC, FL—13; Grissom AFB, IN—26; Holloman AFB, NM—20; Altus AFB, OK—3.

Table 2. Trends in abundance and distribution of *Ae. albopictus* and *Ae. aegypti* from the U.S. Air Force ovitrapping program from 1988 through 1992.

|                            | <i>Ae. albopictus</i> |      |      |      |      | <i>Ae. aegypti</i> |      |      |      |      |
|----------------------------|-----------------------|------|------|------|------|--------------------|------|------|------|------|
|                            | 1988                  | 1989 | 1990 | 1991 | 1992 | 1988               | 1989 | 1990 | 1991 | 1992 |
| No. organizations positive | 10                    | 11   | 12   | 16   | 19   | 12                 | 10   | 4    | 9    | 6    |
| % organizations positive   | 27                    | 29   | 32   | 43   | 68   | 32                 | 26   | 11   | 24   | 21   |
| No. ovipaddles positive    | 56                    | 239  | 118  | 449  | 401  | 91                 | 149  | 11   | 24   | 10   |
| % ovipaddles positive      | 2.7                   | 8.3  | 6.0  | 21.5 | 24.9 | 4.4                | 5.2  | 0.6  | 1.1  | 0.6  |

the abundance of this species has declined dramatically since the establishment of *Ae. albopictus* (McHugh 1991). The single *Ae. aegypti*-positive ovipaddle collected August 19 at Goodfellow AFB, TX, is noteworthy because this base is near the edge of, or just outside, the extreme northwestern range for this species (Darsie and Ward 1981). The few published collection records for the city of San Angelo and Tom Green County (Tinker and Hayes 1959, Morlan and Tinker 1965) indicate that *Ae. aegypti* previously was absent from this area. As was the case with *Ae. albopictus* collected at Goodfellow AFB, the paucity of collections during previous years makes it impossible to determine when or how often *Ae. aegypti* was introduced into the area.

No specimens could be reared from 144 ovipaddles that held eggs tentatively identified as *Aedes* (*Stegomyia*) sp. Most of these eggs had either dried and collapsed or hatched prior to their arrival at Brooks AFB. In previous years, *Aedes* species other than *Ae. triseriatus*, *Ae. albopictus*, and *Ae. aegypti* have been relatively rare; *Aedes epactius* Dyar and Knab was reared from 6 ovipaddles collected from 1988 through 1990 and was absent in 1991 and 1992 rearings. No *Aedes atropalpus* (Coquillett) was reared during 1992. Thus the vast majority of *Aedes* (*Stegomyia*)-positive ovipaddles in the 1992 collections probably were *Ae. albopictus* or *Ae. aegypti*.

Several studies (e.g., Hobbs et al. 1991) have documented a decline in the abundance of *Ae. aegypti* subsequent to the introduction of *Ae. albopictus*. In general, data from the USAF ovitrapping program support this trend. During the years 1988–92, the number of installations positive for *Ae. albopictus* has doubled and the percent of *Ae. albopictus*-positive ovipaddles has increased almost 10-fold (Table 2). During the same period, the distribution and abundance of *Ae. aegypti* have declined markedly. However, *Ae. aegypti* has not been extirpated, even in areas with a high prevalence of *Ae. albopictus*. In Bexar County, TX, for example, *Ae. albopictus* is common, with 30–50%+ of ovipaddles positive for this species during the past few years. *Aedes aegypti* persists in this area, albeit at low levels.

Even in the absence of *Ae. albopictus*, the abundance of *Ae. aegypti* at Seymour Johnson AFB, NC, has declined erratically, with 6.4, 1.7, 0.0, 1.3 and 3.5% positive ovipaddles during 1988–92, respectively. Similarly, Hobbs et al. (1991) reported a marked decline in the abundance of *Ae. aegypti* over several decades prior to the arrival of *Ae. albopictus*. Thus the relative importance of *Ae. albopictus* and other, unrecognized factors in the decline of *Ae. aegypti* remain to be determined.

Thirteen bases submitted ovipaddles positive for *Ae. triseriatus*. No *Aedes hendersoni* Cockerell were reared during 1992.

Forty-six of the ovipaddles were positive for more than one species of *Aedes*. Forty-four were positive for both *Ae. albopictus* and *Ae. triseriatus*. These ovipaddles were submitted by Columbus AFB, MS (16 ovipaddles), Arnold AFS, TN (15), Little Rock AFB, AR (5), Maxwell AFB, AL (3) and Barksdale AFB, LA (2). One ovipaddle collected at Lackland AFB, TX, was positive for both *Ae. albopictus* and *Ae. aegypti*. One ovipaddle collected August 27 at Kelly AFB, TX, was positive for *Ae. albopictus*, *Ae. aegypti* and *Ae. triseriatus*.

D. E. Bowles and M. L. Womack reviewed the manuscript.

#### REFERENCES CITED

- Darsie, R. F., Jr. and R. A. Ward. 1981. Identification and geographical distribution of the mosquitoes of North America, north of Mexico. Mosq. Syst. Suppl. 1.
- Hobbs, J. H., E. A. Hughes and B. H. Eichold II. 1991. Replacement of *Aedes aegypti* by *Aedes albopictus* in Mobile, Alabama. J. Am. Mosq. Control Assoc. 7:488–489.
- Linley, J. R. 1989a. Scanning electron microscopy of the egg of *Aedes* (*Protomacleana*) *triseriatus* (Diptera: Culicidae). J. Med. Entomol. 26:474–478.
- Linley, J. R. 1989b. Comparative fine structure of the eggs of *Aedes albopictus*, *Ae. aegypti*, and *Ae. bahamensis* (Diptera: Culicidae). J. Med. Entomol. 26:510–521.
- McHugh, C. P. 1991. Distributional records from the U.S. Air Force ovitrapping program—1990. J. Am. Mosq. Control Assoc. 7:499–501.

- McHugh, C. P. 1992. Distributional records from the U.S. Air Force ovitrapping program—1991. *J. Am. Mosq. Control Assoc.* 8:198-199.
- McHugh, C. P. and P. A. Hanny. 1990. Records of *Aedes albopictus*, *Ae. aegypti* and *Ae. triseriatus* from the U.S. Air Force ovitrapping program—1989. *J. Am. Mosq. Control Assoc.* 6:549-551.
- McHugh, C. P. and A. M. Vande Berg. 1989. Records of *Aedes albopictus*, *Ae. Aegypti* and *Ae. triseriatus* from the U.S. Air Force ovitrapping program—1988. *J. Am. Mosq. Control Assoc.* 5:440-443.
- Mitchell, C. J., M. L. Niebylski, G. C. Smith, N. Karabatsos, D. Martin, J.-P. Mutebi, G. B. Craig, Jr. and M. J. Mahler. 1992. Isolation of eastern equine encephalitis virus from *Aedes albopictus* in Florida. *Science* 257:526-527.
- Morlan, H. B. and M. E. Tinker. 1965. Distribution of *Aedes aegypti* infestations in the United States. *Am. J. Trop. Med. Hyg.* 14:892-899.
- O'Meara, G. F., A. D. Gettman, L. F. Evans, Jr. and F. D. Scheel. 1992. Invasion of cemeteries in Florida by *Aedes albopictus*. *J. Am. Mosq. Control Assoc.* 8:1-10.
- Tinker, M. E. and G. R. Hayes, Jr. 1959. The 1958 *Aedes aegypti* distribution in the United States. *Mosq. News* 19:73-78.

DTIC QUALITY INSPECTED 5

|                    |                                     |
|--------------------|-------------------------------------|
| Accession For      |                                     |
| NTIS GRA&I         | <input checked="" type="checkbox"/> |
| DTIC TAB           | <input type="checkbox"/>            |
| Unannounced        | <input type="checkbox"/>            |
| Justification      |                                     |
| By                 |                                     |
| Distribution/      |                                     |
| Availability Codes |                                     |
| Dist               | Avail and/or Special                |
| A-1                | 20                                  |